Carbohydrate Research, 227 (1992) xi–xiii
Elsevier Science Publishers B.V., Amsterdam

CONTENTS

Special issue: In Honour of Professor David John Manners

Professor David John Manners vii

Influence of the culture medium on the synthesis of α-D-glucans by *Streptococcus cricetus* AHT
G. J. Walker, V. L. Jacques, E. Fiala-Beer (Surry Hills, N.S.W., Australia), and M. E. Slodki (Peoria, IL, U.S.A.) 1

The cellobiohydrolase I from *Trichoderma reesei* QM 9414: action on cello-oligosaccharides
M. Vršanská and P. Biely (Bratislava, Czechoslovakia) 19

Roles of the aromatic side chains in the binding of substrates, inhibitors, and cyclomalto-oligosaccharides to the glucoamylase from *Aspergillus niger* probed by perturbation difference spectroscopy, chemical modification, and mutagenesis
B. Svensson and M. R. Sierks (Valby, Denmark) 29

Characterization of seven xyloglucan oligosaccharides containing from seventeen to twenty glycosyl residues
M. Hisamatsu, W. S. York, A. G. Darvill, and P. Albersheim (Athens, GA, U.S.A.) . . . 45

The influence of chain length on the hydrodynamic behaviour of amylose
P. Roger and P. Colonna (Nantes, France) 73

Structural changes during development in the amylose and amylopectin fractions (separated by precipitation with concanavalin A) of starches from maize genotypes
S.-H. Yun and N. K. Matheson (Sydney, N.S.W., Australia) 85

Loss of crystalline and molecular order during starch gelatinisation: origin of the enthalpic transition
D. Cooke and M. J. Gidley (Bedford, Gt. Britain) 103

Examination of the structure of amylose by tritium labelling of the reducing terminal
Y. Takeda, N. Maruta, and S. Hizukuri (Kagoshima, Japan) 113

Granule residues and "ghosts" remaining after heating A-type barley-starch granules in water
R. D. M. Prentice, J. R. Stark (Edinburgh, Gt. Britain), and M. J. Gidley (Bedford, Gt. Britain) 121

The structure of four waxy starches related to gelatinization and retrogradation
Y.-C. Shi and P. A. Seib (Manhattan, KS, U.S.A.) 131

The polysaccharides of agricultural lupin seeds
M. T. Al-Kaisey and K. C. B. Wilkie (Old Aberdeen, Gt. Briain) 147

Modification of starch granules by hydrolysis with hydrochloric acid in various alcohols, and the formation of new kinds of limit dextrins
J. D. Fox and J. F. Robyt (Ames, IA, U.S.A.) 163

Characterisation of the (1→4)-α-D-glucan-branching 6-glycosyltransferase by *in vitro* synthesis of branched starch polysaccharides
W. Praznik, G. Rammesmayer, T. Spies (Vienna, Austria), and A. Huber (Graz, Austria) 171

The fine structure of oyster glycogen
M. R. S. Rani, K. Shibanuma, and S. Hizukuri (Kagoshima, Japan) 183

Monitoring polysaccharide synthesis in *Candida albicans*
J. A. Brown (Edinburgh, Gt. Britain) and B. J. Catley (Dunedin, New Zealand) 195

A study of the heat capacity of starch/water mixtures
T. R. Noel and S. G. Ring (Norwich, Gt. Britain) 203

Action patterns of amylolytic enzymes as determined by the [1-^{14}C]malto-oligosaccharide mapping method
J. H. Pazur and N. T. Marchetti (University Park, PA, U.S.A.) 215

Characterization of ADPglucose pyrophosphorylase from a starch-deficient mutant of *Arabidopsis thaliana* (L.)
L. Li and J. Preiss (East Lansing, MA, U.S.A.) 227

(1→4)-α-D-Glucan synthesis by a chloroplastic phosphorylase isolated from spinach leaves is independent of added primer
M. N. Sivak (East Lansing, MI, U.S.A.) 241

Measurement of the content of limit-dextrinase in cereal flours
B. V. McCleary (North Rocks, N.S.W., Australia) 257

A method for the study of the enzymic hydrolysis of starch granules
E. Bertoft and R. Manelius (Turku, Finland) 269

Effect of modification on the tryptophan residues of cyclodextrin glucanotransferase with *N*-bromosuccinimide on the enzyme-catalysed hydrolysis (cleavage) of soluble starch and cyclomaltohexaose
M. Ohnishi, U. Ota, M. Abe, B. Tonomura (Kyoto, Japan), and M. Kubota (Okayama, Japan) 285

The cellulase of the anaerobic bacterium *Clostridium thermocellum*: isolation, dissociation, and reassociation of the cellulosome
K. M. Bhat and T. M. Wood (Aberdeen, Gt. Britain) 293

The action of germinated barley alpha-amylases on linear maltodextrins
A. W. MacGregor, J. E. Morgan, and E. A. MacGregor (Winnipeg, Man., Canada) . . . 301

Structural characterization of an arabinoxylan–rhamnogalacturonan complex from cell walls of *Zea* shoots
Y. Kato (Hirosaki, Japan) and D. J. Nevins (Davis, CA, U.S.A.) 315

The substrate specificity of isoamylase and the preparation of apo-glycogenin
J. Lomako, W. M. Lomako, and W. J. Whelan (Miami, FL, U.S.A.) 331

Molecular and metabolic aspects of lysosomal glycogen
R. Geddes, P. Jeyarathan, and J. A. Taylor (Auckland, New Zealand) 339

Molecular modelling of acarviosine, the pseudo-disaccharide moiety of acarbose, and other inhibitors of alpha-amylases
E. Raimbaud, A. Buleon, and S. Perez (Nantes, France) 351

Notes

The detection and quantification of apiose by capillary gas chromatography of its alditol acetates
P. J. Harris (Parkville, Vict., Australia), M. Sadek, R. T. C. Brownlee (Bundoora, Vict., Australia), A. B. Blakeney (Yanco, N.S.W., Australia), J. Webster, and B. A. Stone (Bundoora, Vict., Australia) 365

Conformation of 3-*O*-*β*-D-galactopyranosyl-L-arabinose and a comparison with its α-linked isomer
D. C. Vogt, G. E. Jackson, and A. M. Stephen (Rondebosch, South Africa) 371

A re-investigation of the borohydride reduction of carbohydrates
R. J. Sturgeon (Edinburgh, Gt. Britain) 375

The action of isoamylase on the surface of starch granules
A. Lynn and J. R. Stark (Edinburgh, Gt. Britain) 379

Heterogeneity of the glucoamylase components of the raw-starch-digesting amylase from *Chalara paradoxa*
M. Monma and K. Kainuma (Tsukuba, Japan) 385

Preliminary communication
The amylase of *Pseudomonas stutzeri* as a probe of the structure of amylopectin
P. Finch and D. W. Sebesta (Egham, Gt. Britain) c1

Author index c5

Subject index c7